

HARDY-TYPE INEQUALITIES VIA AUXILIARY SEQUENCES

PENG GAO

ABSTRACT. We prove some Hardy-type inequalities via an approach that involves constructing auxiliary sequences.

1. INTRODUCTION

Suppose throughout that $p \neq 0$, $\frac{1}{p} + \frac{1}{q} = 1$. Let l^p be the Banach space of all complex sequences $\mathbf{a} = (a_n)_{n \geq 1}$ with norm

$$\|\mathbf{a}\| := \left(\sum_{n=1}^{\infty} |a_n|^p \right)^{1/p} < \infty.$$

The celebrated Hardy's inequality ([5, Theorem 326]) asserts that for $p > 1$,

$$(1.1) \quad \sum_{n=1}^{\infty} \left| \frac{1}{n} \sum_{k=1}^n a_k \right|^p \leq \left(\frac{p}{p-1} \right)^p \sum_{k=1}^{\infty} |a_k|^p.$$

Hardy's inequality can be regarded as a special case of the following inequality:

$$\sum_{j=1}^{\infty} \left| \sum_{k=1}^{\infty} c_{j,k} a_k \right|^p \leq U \sum_{k=1}^{\infty} |a_k|^p,$$

in which $C = (c_{j,k})$ and the parameter p are assumed fixed ($p > 1$), and the estimate is to hold for all complex sequences $\mathbf{a}$. The l^p operator norm of C is then defined as the p -th root of the smallest value of the constant U :

$$\|C\|_{p,p} = U^{\frac{1}{p}}.$$

Hardy's inequality thus asserts that the Cesàro matrix operator C , given by $c_{j,k} = 1/j, k \leq j$ and 0 otherwise, is bounded on l^p and has norm $\leq p/(p-1)$. (The norm is in fact $p/(p-1)$.)

We say a matrix A is a summability matrix if its entries satisfy: $a_{j,k} \geq 0$, $a_{j,k} = 0$ for $k > j$ and $\sum_{k=1}^j a_{j,k} = 1$. We say a summability matrix A is a weighted mean matrix if its entries satisfy:

$$a_{j,k} = \lambda_k / \Lambda_j, \quad 1 \leq k \leq j; \quad \Lambda_j = \sum_{i=1}^j \lambda_i, \quad \lambda_i \geq 0, \quad \lambda_1 > 0.$$

Hardy's inequality (1.1) now motivates one to determine the l^p operator norm of an arbitrary summability matrix A . For examples, the following two inequalities were claimed to hold by Bennett ([1, p. 40-41]; see also [2, p. 407]):

$$(1.2) \quad \sum_{n=1}^{\infty} \left| \frac{1}{n^\alpha} \sum_{i=1}^n (i^\alpha - (i-1)^\alpha) a_i \right|^p \leq \left(\frac{\alpha p}{\alpha p - 1} \right)^p \sum_{n=1}^{\infty} |a_n|^p,$$

$$(1.3) \quad \sum_{n=1}^{\infty} \left| \frac{1}{\sum_{i=1}^n i^{\alpha-1}} \sum_{i=1}^n i^{\alpha-1} a_i \right|^p \leq \left(\frac{\alpha p}{\alpha p - 1} \right)^p \sum_{n=1}^{\infty} |a_n|^p,$$

Date: May 24, 2007.

2000 *Mathematics Subject Classification*. Primary 26D15.

Key words and phrases. Hardy's inequality.

whenever $\alpha > 0, p > 1, \alpha p > 1$.

No proofs of the above two inequalities were supplied in [1]-[2] and recently, the author [4] and Bennett himself [3] proved inequalities (1.2) for $p > 1, \alpha \geq 1, \alpha p > 1$ and (1.3) for $p > 1, \alpha \geq 2$ or $0 < \alpha \leq 1, \alpha p > 1$ independently.

We point out here that Bennett in fact was able to prove (1.2) for $p \geq 1, \alpha > 0, \alpha p > 1$ (see [3, Theorem 1] with $\beta = 1$ there) which now leaves the case $p > 1, 1 < \alpha < 2$ of inequality (1.3) the only case open to us. For this, Bennett expects inequality (1.3) to hold for $1 + 1/p < \alpha < 2$ (see page 830 of [3]) and as a support, Bennett [3, Theorem 18] has shown that inequality (1.3) holds for $\alpha = 1 + 1/p, p \geq 1$.

In this paper, we will study inequality (1.3) using a method of Knopp [6] which involves constructing auxiliary sequences. We will partially resolve the remaining case $p > 1, 1 < \alpha < 2$ of inequality (1.3) by proving in Section 2 the following:

Theorem 1.1. *Inequality (1.3) holds for $p \geq 2, 1 \leq \alpha \leq 1 + 1/p$ or $1 < p \leq 4/3, 1 + 1/p \leq \alpha \leq 2$.*

We shall leave the explanation of Knopp's approach in detail in Section 2 by pointing out here that it can be applied to prove other types of inequalities similar to that of Hardy's. As an example, we note that Theorem 359 of [5] states:

Theorem 1.2. *For $0 < p < 1$ and $a_n \geq 0$,*

$$\sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{k=n}^{\infty} a_k \right)^p \geq p^p \sum_{n=1}^{\infty} a_n^p.$$

The constant p^p in Theorem 1.2 is not best possible and this was fixed by Levin and Stečkin [7, Theorem 61] for $0 < p \leq 1/3$ in the following

Theorem 1.3. *For $0 < p \leq 1/3$ and $a_n \geq 0$,*

$$\sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{k=n}^{\infty} a_k \right)^p \geq \left(\frac{p}{1-p} \right)^p \sum_{n=1}^{\infty} a_n^p.$$

We shall give another proof of this result in Section 3 using Knopp's approach. We point out here for each $1/3 < p < 1$, Levin and Stečkin also gave a better constant than the one p^p given in Theorem 1.2. For example, when $p = 1/2$, they gave $\sqrt{3}/2$ instead of $1/\sqrt{2}$. In Section 4, we shall consider an approach of Redheffer [8] by showing first that this approach can be regarded as essentially the approach of Knopp when treating Hardy-type inequalities. We then use Redheffer's method to prove the following

Theorem 1.4. *For $a_n \geq 0$,*

$$\sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{k=n}^{\infty} a_k \right)^{1/2} \geq 0.8967 \sum_{n=1}^{\infty} a_n^{1/2}.$$

This improves the result of Levin and Stečkin mentioned above. It is also pointed out in Section 4 that the same method can be used to establish the result in Theorem 1.3 for p slightly bigger than $1/3$.

In our proofs of Theorems 1.1-1.2, certain auxiliary sequences are constructed and there can be many ways to construct such sequences. In Section 5, we give an example regarding these possibilities by answering a question of Bennett.

2. PROOF OF THEOREM 1.1

We begin this section by explaining Knopp's idea [6] on proving Hardy's inequality (1.1). In fact, we will explain this more generally for the case involving weighted mean matrices. For real

numbers $\lambda_1 > 0, \lambda_i \geq 0, i \geq 2$, we write $\Lambda_n = \sum_{i=1}^n \lambda_i$ and we are looking for a positive constant U such that

$$(2.1) \quad \sum_{n=1}^{\infty} \left| \frac{1}{\Lambda_n} \sum_{k=1}^n \lambda_k a_k \right|^p \leq U \sum_{k=1}^{\infty} |a_k|^p$$

holds for all complex sequences $\mathbf{a}$ with $p > 1$ being fixed. Knopp's idea is to find an auxiliary sequence $\mathbf{w} = \{w_i\}_{i=1}^{\infty}$ of positive terms such that by Hölder's inequality,

$$\begin{aligned} \left(\sum_{k=1}^n \lambda_k |a_k| \right)^p &= \left(\sum_{k=1}^n \lambda_k |a_k| w_k^{-\frac{1}{p^*}} \cdot w_k^{\frac{1}{p^*}} \right)^p \\ &\leq \left(\sum_{k=1}^n \lambda_k^p |a_k|^p w_k^{-(p-1)} \right) \left(\sum_{j=1}^n w_j \right)^{p-1} \end{aligned}$$

so that

$$\begin{aligned} \sum_{n=1}^{\infty} \left| \frac{1}{\Lambda_n} \sum_{k=1}^n \lambda_k a_k \right|^p &\leq \sum_{n=1}^{\infty} \frac{1}{\Lambda_n^p} \left(\sum_{k=1}^n \lambda_k^p |a_k|^p w_k^{-(p-1)} \right) \left(\sum_{j=1}^n w_j \right)^{p-1} \\ &= \sum_{k=1}^{\infty} w_k^{-(p-1)} \lambda_k^p \left(\sum_{n=k}^{\infty} \frac{1}{\Lambda_n^p} \left(\sum_{j=1}^n w_j \right)^{p-1} \right) |a_k|^p. \end{aligned}$$

Suppose now one can find for each $p > 1$ a positive constant U , a sequence $\mathbf{w}$ of positive terms with w_n^{p-1}/λ_n^p decreasing to 0, such that for any integer $n \geq 1$,

$$(2.2) \quad (w_1 + \cdots + w_n)^{p-1} < U \Lambda_n^p \left(\frac{w_n^{p-1}}{\lambda_n^p} - \frac{w_{n+1}^{p-1}}{\lambda_{n+1}^p} \right),$$

then it is easy to see that inequality (2.1) follows from this. When $\lambda_n = 1$ for all n , Knopp's choice for $\mathbf{w}$ is given by $w_n = \binom{n-1-1/p}{n-1}$ and one can show that (2.2) holds in this case with $U = (p^*)^p$ and Hardy's inequality (1.1) follows from this.

We now want to apply Knopp's approach to prove Theorem 1.1. For this, we replace $\alpha - 1$ by α and rewrite (1.3) as

$$(2.3) \quad \sum_{n=1}^{\infty} \left| \frac{1}{\sum_{i=1}^n i^\alpha} \sum_{i=1}^n i^\alpha a_i \right|^p \leq \left(\frac{(\alpha+1)p}{(\alpha+1)p-1} \right)^p \sum_{n=1}^{\infty} |a_n|^p.$$

Note that we are interested in the case $0 \leq \alpha \leq 1$ here. From our discussions above, we are looking for a sequence $\mathbf{w}$ of positive terms with w_n^{p-1}/λ_n^p decreasing to 0, such that for any integer $n \geq 1$,

$$(2.4) \quad (w_1 + \cdots + w_n)^{p-1} < \left(\frac{(\alpha+1)p}{(\alpha+1)p-1} \right)^p \left(\sum_{i=1}^n i^\alpha \right)^p \left(\frac{w_n^{p-1}}{n^{\alpha p}} - \frac{w_{n+1}^{p-1}}{(n+1)^{\alpha p}} \right).$$

Following Knopp's choice, we define a sequence $\mathbf{w}$ such that

$$(2.5) \quad w_{n+1} = \frac{n + \alpha - 1/p}{n} w_n, \quad n \geq 1.$$

Note that the above sequence is uniquely determined for any given positive w_1 and therefore we may assume $w_1 = 1$ here. We note further that we need $\alpha > -1/p^*$ in order for $w_n > 0$ for all n and we also point out that it is easy to show by induction that

$$(2.6) \quad \sum_{i=1}^n w_i = \frac{n + \alpha - 1/p}{1 + \alpha - 1/p} w_n.$$

Moreover, one can easily check that

$$\frac{w_n^{p-1}}{n^{\alpha p}} = O(n^{-\alpha-1/p^*}),$$

so that w_n^{p-1}/λ_n^p decreases to 0 as n approaches infinity as long as $\alpha > -1/p^*$.

Now we need a lemma on sums of powers, which is due to Levin and Stečkin [7, Lemma 1, 2, p.18]:

Lemma 2.1. *For an integer $n \geq 1$,*

$$(2.7) \quad \sum_{i=1}^n i^r \geq \frac{1}{r+1} n(n+1)^r, \quad 0 \leq r \leq 1,$$

$$(2.8) \quad \sum_{i=1}^n i^r \geq \frac{r}{r+1} \frac{n^r(n+1)^r}{(n+1)^r - n^r}, \quad r \geq 1.$$

Inequality (2.8) reverses when $-1 < r \leq 1$.

We note here only the case $r \geq 0$ for (2.8) was proved in [7] but one checks easily that the proof extends to the case $r > -1$.

As we are interested in $0 \leq \alpha \leq 1$ here, we can now combine (2.5)-(2.7) to deduce that inequality (2.4) will follow from

$$\left(1 + \frac{\alpha - 1/p}{n}\right)^{p-1} < \frac{n}{1 + \alpha - 1/p} \left(\left(1 + \frac{1}{n}\right)^{\alpha p} - \left(1 + \frac{\alpha - 1/p}{n}\right)^{p-1} \right).$$

We can simplify the above inequality further by recasting it as

$$(2.9) \quad \left(1 + \frac{\alpha + 1/p^*}{n}\right)^{1/p} \left(1 + \frac{\alpha - 1/p}{n}\right)^{1/p^*} < \left(1 + \frac{1}{n}\right)^{\alpha}.$$

Now we define for fixed $n \geq 1, p > 1$,

$$f(x) = x \ln(1 + 1/n) - \frac{1}{p} \ln\left(1 + \frac{x + 1/p^*}{n}\right) - \frac{1}{p^*} \ln\left(1 + \frac{x - 1/p}{n}\right).$$

It is easy to see here that inequality (2.9) is equivalent to $f(\alpha) > 0$. It is also easy to see that $f(x)$ is a convex function of x for $0 \leq x \leq 1$ and that $f(1/p) = 0$. It follows from this that if $f'(1/p) \leq 0$ then $f(x) > 0$ for $0 \leq x < 1/p$ and if $f'(1/p) \geq 0$ then $f(x) > 0$ for $1/p < x \leq 1$. We have

$$f'(1/p) = \ln(1 + 1/n) - \frac{1}{n} + \frac{1}{pn(n+1)}.$$

We now use Taylor expansion to conclude for $x > 0$,

$$(2.10) \quad x - x^2/2 < \ln(1 + x) < x - x^2/2 + x^3/3.$$

It follows from this that for $p \geq 2, n \geq 2$,

$$f'(1/p) < -\frac{1}{2n^2} + \frac{1}{3n^3} + \frac{1}{pn(n+1)} \leq -\frac{1}{2n^2} + \frac{1}{3n^3} + \frac{1}{2n(n+1)} = \frac{1}{3n^3} - \frac{1}{2n^2(n+1)} \leq 0.$$

and for $n = 1$,

$$f'(1/p) = \ln 2 - 1 + \frac{1}{2p} \leq \ln 2 - 1 + \frac{1}{4} < 0,$$

It's also easy to check that for $1 < p \leq 4/3, n = 1$,

$$f'(1/p) = \ln 2 - 1 + \frac{1}{2p} > 0.$$

For $n \geq 2, 1 < p \leq 4/3$, by using the first inequality of (2.10) we get

$$f'(1/p) > -\frac{1}{2n^2} + \frac{1}{pn(n+1)} \geq 0.$$

This now enables us to conclude the proof of Theorem 1.1.

3. ANOTHER PROOF OF THEOREM 1.3

We use the idea of Levin and Stečkin in the proof of Theorem 62 in [7] to find an auxiliary sequence $\mathbf{w} = \{w_i\}_{i=1}^{\infty}$ of positive terms so that for any finite summation from $n = 1$ to N with $N \geq 1$, we have

$$\sum_{n=1}^N a_n^p = \sum_{n=1}^N \frac{a_n^p}{\sum_{i=1}^n w_i} \sum_{k=1}^n w_k = \sum_{n=1}^N w_n \sum_{k=n}^N \frac{a_k^p}{\sum_{i=1}^k w_i}.$$

On letting $N \rightarrow \infty$, we then have

$$\sum_{n=1}^{\infty} a_n^p = \sum_{n=1}^{\infty} w_n \sum_{k=n}^{\infty} \frac{a_k^p}{\sum_{i=1}^k w_i}.$$

By Hölder's inequality, we have

$$\sum_{k=n}^{\infty} \frac{a_k^p}{\sum_{i=1}^k w_i} \leq \left(\sum_{k=n}^{\infty} \left(\sum_{i=1}^k w_i \right)^{-1/(1-p)} \right)^{1-p} \left(\sum_{k=n}^{\infty} a_k^p \right)^p.$$

Suppose now one can find a sequence $\mathbf{w}$ of positive terms with $w_n^{-1/(1-p)} n^{-p/(1-p)}$ decreasing to 0 for each $0 < p \leq 1/3$, such that for any integer $n \geq 1$,

$$(3.1) \quad (w_1 + \cdots + w_n)^{-1/(1-p)} \leq \left(\frac{1-p}{p} \right)^{p/(1-p)} \left(\frac{w_n^{-1/(1-p)}}{n^{p/(1-p)}} - \frac{w_{n+1}^{-1/(1-p)}}{(n+1)^{p/(1-p)}} \right),$$

then it is easy to see that Theorem 1.3 follows from this.

We now define our sequence $\mathbf{w}$ to be

$$(3.2) \quad w_{n+1} = \frac{n+1/p-2}{n} w_n, \quad n \geq 1.$$

Note that the above sequence is uniquely determined for any given positive w_1 and therefore we may assume $w_1 = 1$ here. We note further that $w_n > 0$ for all n as $0 < p \leq 1/3$ and it is easy to show by induction that

$$(3.3) \quad \sum_{i=1}^n w_i = \frac{n+1/p-2}{1/p-1} w_n.$$

Moreover, one can easily check that

$$\frac{w_n^{-1/(1-p)}}{n^{p/(1-p)}} = O(n^{-(1-p)/p}),$$

so that $w_n^{-1/(1-p)} n^{-p/(1-p)}$ decreases to 0 as n approaches infinity.

We now combine (3.2)-(3.3) to recast inequality (3.1) as

$$(n+1/p-2)^{-1/(1-p)} \leq \frac{p}{1-p} \left(n^{-p/(1-p)} - (n+1)^{-p/(1-p)} n^{1/(1-p)} (n+1/p-2)^{-1/(1-p)} \right).$$

We further rewrite the above inequality as

$$\begin{aligned} \frac{1-p}{p} &\leq n^{-p/(1-p)}(n+1/p-2)^{1/(1-p)} - (n+1)^{-p/(1-p)}n^{1/(1-p)} \\ &= n\left(\left(1+\frac{1/p-2}{n}\right)^{1/(1-p)} - \left(1+\frac{1}{n}\right)^{-p/(1-p)}\right). \end{aligned}$$

It is easy to see that the above inequality follows from $f(1/n) \geq 0$ where we define for $x \geq 0$,

$$f(x) = \left(1 + (1/p - 2)x\right)^{1/(1-p)} - \left(1 + x\right)^{-p/(1-p)} - \frac{1-p}{p}x.$$

We now prove that $f(x) \geq 0$ for $x \geq 0$ for $0 < p \leq 1/3$ and this will conclude the proof of Theorem 1.3. We note that

$$\begin{aligned} f'(x) &= \frac{1/p-2}{1-p} \left(1 + (1/p-2)x\right)^{p/(1-p)} + \frac{p}{1-p} \left(1+x\right)^{-p/(1-p)-1} - \frac{1-p}{p}, \\ f''(x) &= \frac{p(1/p-2)^2}{(1-p)^2} \left(1 + (1/p-2)x\right)^{p/(1-p)-1} - \frac{p}{(1-p)^2} \left(1+x\right)^{-p/(1-p)-2}. \end{aligned}$$

We now define for $x \geq 0$,

$$g(x) = (1/p-2)^{2(1-p)/(1-2p)}(1+x)^{(2-p)/(1-2p)} - (1 + (1/p-2)x).$$

It is easy to see that $g(x) \geq 0$ implies $f''(x) \geq 0$. Note that $(2-p)/(1-2p) \geq 1$ so that

$$\begin{aligned} g'(x) &= (1/p-2)^{2(1-p)/(1-2p)}(2-p)/(1-2p)(1+x)^{(2-p)/(1-2p)-1} - (1/p-2) \\ &\geq (1/p-2)^{2(1-p)/(1-2p)} - (1/p-2) \geq 0, \end{aligned}$$

where the last inequality above follows from $2(1-p)/(1-2p) \geq 1$ and $0 < p \leq 1/3$ so that $1/p-2 \geq 1$. It follows from this that $f''(x) \geq 0$ and as one checks easily that $f'(0) = 0$, which implies $f'(x) \geq 0$ so that $f(x) \geq f(0) = 0$ which is just what we want to prove.

4. REDHEFFER'S APPROACH AND PROOF OF THEOREM 1.4

Redheffer's approach in [8] of Hardy-type inequalities via his "recurrent inequalities" can be put into the following form:

Lemma 4.1 ([4, Lemma 2.4]). *Let $\{\lambda_i\}_{i \geq 1}^\infty, \{a_i\}_{i \geq 1}^\infty$ be two sequences of positive real numbers and let $S_n = \sum_{i=1}^n \lambda_i a_i$. Let $0 \neq p < 1$ be fixed and let $\{\mu_i\}_{i \geq 1}^\infty, \{\eta_i\}_{i \geq 1}^\infty$ be two positive sequences of real numbers such that $\mu_i \leq \eta_i$ for $0 < p < 1$ and $\mu_i \geq \eta_i$ for $p < 0$, then for $n \geq 2$,*

$$(4.1) \quad \sum_{i=2}^{n-1} \left(\mu_i - (\mu_{i+1}^q - \eta_{i+1}^q)^{1/q} \right) S_i^{1/p} + \mu_n S_n^{1/p} \leq (\mu_2^q - \eta_2^q)^{1/q} \lambda_1^{1/p} a_1^{1/p} + \sum_{i=2}^n \eta_i \lambda_i^{1/p} a_i^{1/p}.$$

We consider the case $0 < p < 1$ in the above lemma and we set $\eta_i = \lambda_i^{-1/p}$ together with a change of variables: $\mu_i \mapsto \mu_i \eta_i$ to rewrite (4.1) as

$$\sum_{i=2}^{n-1} \left(\frac{\mu_i}{\lambda_i^{1/p}} - \frac{(\mu_{i+1}^q - 1)^{1/q}}{\lambda_{i+1}^{1/p}} \right) S_i^{1/p} + \frac{\mu_n}{\lambda_n^{1/p}} S_n^{1/p} \leq (\mu_2^q - 1)^{1/q} \frac{\lambda_1^{1/p}}{\lambda_2^{1/p}} a_1^{1/p} + \sum_{i=2}^n a_i^{1/p}.$$

We now set $\mu_i^q - 1 = \nu_i$ and make a further change of variables: $p \mapsto 1/p$ to write the above inequality as:

$$(4.2) \quad \sum_{i=2}^{n-1} \left(\frac{(1+\nu_i)^{-(p-1)}}{\lambda_i^p} - \frac{\nu_{i+1}^{-(p-1)}}{\lambda_{i+1}^p} \right) S_i^p + \frac{(1+\nu_n)^{-(p-1)}}{\lambda_n^p} S_n^p \leq \nu_2^{-(p-1)} \frac{\lambda_1^p}{\lambda_2^p} a_1^p + \sum_{i=2}^n a_i^p.$$

Now if we set for $i \geq 2$,

$$\nu_i = \frac{\sum_{j=1}^{i-1} w_j}{w_i},$$

we can rewrite inequality (4.2) as

$$(4.3) \quad \begin{aligned} & \sum_{i=2}^{n-1} \left(\sum_{j=1}^i w_j \right)^{-(p-1)} \left(\frac{w_i^{p-1}}{\lambda_i^p} - \frac{w_{i+1}^{p-1}}{\lambda_{i+1}^p} \right) \Lambda_i^p A_i^p + \left(\sum_{j=1}^n w_j \right)^{-(p-1)} \frac{w_n^{p-1}}{\lambda_n^p} \Lambda_n^p A_n^p \\ & \leq \frac{w_2^{p-1}}{w_1^{p-1}} \frac{\lambda_1^p}{\lambda_2^p} a_1^p + \sum_{i=2}^n a_i^p, \end{aligned}$$

where

$$\Lambda_n = \sum_{i=1}^n \lambda_i, \quad A_n = \frac{S_n}{\Lambda_n}, \quad n \geq 1.$$

Suppose now we can find a sequence $\mathbf{w} = \{w_i\}_{i=1}^\infty$ of positive terms such that inequality (2.2) holds for all $n \geq 1$. Then inequality (4.3) implies

$$(4.4) \quad \frac{1}{U} \sum_{i=1}^n A_i^p \leq \left(\frac{1}{U} + \frac{w_2^{p-1}}{w_1^{p-1}} \frac{\lambda_1^p}{\lambda_2^p} \right) a_1^p + \sum_{i=2}^n a_i^p \leq \sum_{i=1}^n a_i^p,$$

where the last inequality above follows from the case $n = 1$ of inequality (2.2), which implies

$$\frac{1}{U} < 1 - \frac{w_2^{p-1}}{w_1^{p-1}} \frac{\lambda_1^p}{\lambda_2^p}.$$

Thus we have seen that on letting $n \rightarrow +\infty$, inequality (4.4) gives back inequality (2.1). Hence Redheffer's approach can be regarded as essentially Knopp's approach when treating Hardy-type inequalities. The only difference is that one no longer requires that w_n^{p-1}/λ_n^p decreases to 0 when selecting the sequence $\mathbf{w}$ in Redheffer's approach.

Now we state a lemma similar to Lemma 4.1:

Lemma 4.2. *Let $\{\lambda_i\}_{i \geq 1}^\infty, \{a_i\}_{i \geq 1}^\infty$ be two sequences of positive real numbers and suppose $\sum_{i=1}^\infty \lambda_i a_i$ converges. Let $S_n = \sum_{i=n}^\infty \lambda_i a_i$ and let $0 < p < 1$ be fixed. Let $\{\mu_i\}_{i \geq 1}^\infty, \{\eta_i\}_{i \geq 1}^\infty$ be two positive sequences of real numbers such that $\mu_i \geq \eta_i$, then for $n \geq 2$,*

$$(4.5) \quad \mu_1 S_1^p + \sum_{i=2}^n \left(\mu_i - (\mu_{i-1}^{\frac{1}{1-p}} - \eta_{i-1}^{\frac{1}{1-p}})^{1-p} \right) S_i^p - (\mu_n^{\frac{1}{1-p}} - \eta_n^{\frac{1}{1-p}})^{1-p} S_{n+1}^p \geq \sum_{i=1}^n \eta_i \lambda_i^p a_i^p.$$

Proof. We note for $k \geq 2$,

$$(4.6) \quad \mu_k S_k^p - \eta_k \lambda_k^p a_k^p = S_{k+1}^p (\mu_k (1+t)^p - \eta_k t^p) \geq (\mu_k^{\frac{1}{1-p}} - \eta_k^{\frac{1}{1-p}})^{1-p} S_{k+1}^p,$$

with $t = \lambda_k a_k / S_{k+1}$. The lemma then follows by adding (4.6) for $1 \leq k \leq n$ together. $\square$

We set $\eta_i = \lambda_i^{-p}$ together with a change of variables: $\mu_i \mapsto \mu_i \eta_i$ to rewrite (4.5) as

$$\frac{\mu_1}{\lambda_1^p} S_1^p + \sum_{i=2}^n \left(\frac{\mu_i}{\lambda_i^p} - \frac{(\mu_{i-1}^{\frac{1}{1-p}} - 1)^{1-p}}{\lambda_{i-1}^p} \right) S_i^p - \frac{(\mu_n^{\frac{1}{1-p}} - 1)^{1-p}}{\lambda_n^p} S_{n+1}^p \geq \sum_{i=1}^n a_i^p.$$

We now set $\mu_i^{\frac{1}{1-p}} - 1 = \nu_i$ and write the above inequality as:

$$\frac{(1+\nu_1)^{1-p}}{\lambda_1^p} S_1^p + \sum_{i=2}^n \left(\frac{(1+\nu_i)^{1-p}}{\lambda_i^p} - \frac{\nu_{i-1}^{1-p}}{\lambda_{i-1}^p} \right) S_i^p - \frac{\nu_n^{1-p}}{\lambda_n^p} S_{n+1}^p \geq \sum_{i=1}^n a_i^p.$$

From now on we consider the case $\lambda_i = 1$ for all i in the above inequality and we set for $n \geq 1$,

$$\nu_n = \frac{n - \beta}{c},$$

with $\beta \leq 1, c \geq \beta$ here. We want to choose c, β such that the following inequality holds for $n \geq 2$:

$$(4.7) \quad \max \left((1 + c - \beta)^{1-p}, n^p \left((n + c - \beta)^{1-p} - (n - 1 - \beta)^{1-p} \right) \right) \leq c^{1-p} k(p),$$

where $k(p)$ is a constant depending only on p and we want $k(p)$ to be as small as possible. For this purpose, we further assume that $k(p)$ satisfies:

$$(4.8) \quad (1 - p)(1 + c) < c^{1-p} k(p),$$

and define for $0 \leq x \leq 1/2$,

$$f(x) = (1 + (c - \beta)x)^{1-p} - (1 - (1 + \beta)x)^{1-p} - c^{1-p} k(p)x,$$

and note that with our assumption on $k(p)$, $f'(0) < 0$. Note also that

$$f''(x) = p(1 - p)(1 + \beta)^2(1 - (1 + \beta)x)^{-p-1} - p(1 - p)(c - \beta)^2(1 + (c - \beta)x)^{-p-1}.$$

It follows from this that when $1 + \beta \geq c - \beta$ then $f''(x) \geq 0$ for $0 \leq x \leq 1/2$. Otherwise we note that $f''(x) = 0$ can have at most one root in $(0, 1/2)$ and $f''(0) < 0$. The above implies that for $0 \leq x \leq 1/2$, $f(x) \leq \min(f(0), f(1/2)) = \min(0, f(1/2))$. We deduce from our discussion above on setting $x = 1/n$ in $f(x)$ that in order for inequality (4.7) to hold, it suffices to check the case $n = 2$, namely,

$$(4.9) \quad \max \left((1 + c - \beta)^{1-p}, 2^p \left((2 + c - \beta)^{1-p} - (1 - \beta)^{1-p} \right) \right) \leq c^{1-p} k(p),$$

provided we assume (4.8).

We now look at the case $p = 1/2$ and in this case we choose c, β so that the following holds:

$$\left(1 + \frac{1 - \beta}{c}\right)^{1/2} = 2^{1/2} \left(\left(1 + \frac{2 - \beta}{c}\right)^{1/2} - \left(\frac{1 - \beta}{c}\right)^{1/2} \right).$$

On setting $x = (1 - \beta)/c, c' = 1/c$, we can rewrite the above equation as

$$(1 + x)^{1/2} = 2^{1/2} \left((1 + c' + x)^{1/2} - x^{1/2} \right).$$

Solving the above equation yields:

$$x = \frac{1 - \beta}{c} = \frac{\sqrt{(10 + 4c')^2 + 28(1 + 2c')^2} - (10 + 4c')}{14}.$$

To prove Theorem 1.4, we take $c = 5/2$ here, then $x \approx 0.2435$ with $\beta \approx 0.3912$ and $k(1/2) \approx 1.1151 < 1.1152$. We take $k(p) = 1.1152$ here and one can also check that inequality (4.8) holds in this case. As $1/1.1152 > 0.8967$, Theorem 1.4 now follows.

Now we consider other values of p 's. For this, we choose $c = 1/p - 1, k(p) = c^p$ so that inequality (4.8) becomes an equality. Because of this, we need to assume that

$$(4.10) \quad \beta < \frac{1}{2p} - 1,$$

so that $f''(0) < 0$ is satisfied for the function $f(x)$ defined above and our argument goes through as well in the above discussions to ensure that when inequality (4.9) holds, inequality (4.7) also holds. For the case $p = 1/3$, it is easy to check that on taking $\beta = 3 - 2\sqrt{2}$, both inequalities (4.9) and (4.10) are satisfied. This implies Theorem 1.3 for the case $p = 1/3$. In view of this, one sees that it is possible to prove the result in Theorem 1.3 for p beyond $1/3$. For example, on taking $p = 0.34, \beta = 0.21$, calculations shows both inequalities (4.10) and (4.9) are satisfied and hence Theorem 1.3 holds for $p = 0.34$.

5. ANOTHER LOOK AT INEQUALITY (1.3)

In this section we return to the consideration of inequality (1.3) via our approach in Section 2, which boils down to a construction of a sequence $\mathbf{w}$ of positive terms with w_n^{p-1}/λ_n^p decreasing to 0, such that for any integer $n \geq 1$, inequality (2.4) is satisfied. Certainly here the choice for $\mathbf{w}$ may not be unique and in fact in the case $\alpha = 0$, Bennett asked in [1] (see the paragraph below Lemma 4.11) for other sequences, not multiples of Knopp's, that satisfy (2.4). He also mentioned that the obvious choice, $w_n = n^{-1/p}$, does not work.

We point out here even though the choice $w_n = n^{-1/p}$ does not satisfy (2.4) when $\alpha = 0$ for all $p > 1$, as one can see by considering inequality (2.4) for the case $n = 1$ with $p \rightarrow 1^+$, it nevertheless works for $p \geq 3$, which we now show by first rewriting (2.4) in our case as

$$(5.1) \quad \left(\sum_{i=1}^n i^{-1/p} \right)^{p-1} < \left(\frac{p}{p-1} \right)^p n^p \left(n^{-(p-1)/p} - (n+1)^{-(p-1)/p} \right).$$

We note that the case $n = 1$ of (5.1) follows from the case $\alpha = 0$ of the following inequality,

$$(5.2) \quad 1 - 2^{-(p-1)/p-\alpha} > \left(1 - \frac{1}{(\alpha+1)p} \right)^p, \quad 0 \leq \alpha \leq 1/p.$$

To show (5.2), we see by Taylor expansion, that for $p \geq 2, x < 0$,

$$(1+x)^p < 1 + px + \frac{p(p-1)x^2}{2}.$$

Apply the above inequality with $x = -1/(\alpha p + p)$, we obtain for $p \geq 3$,

$$\left(1 - \frac{1}{(\alpha+1)p} \right)^p < 1 - \frac{1}{(\alpha+1)} + \frac{(p-1)}{2(\alpha+1)^2 p}.$$

Hence inequality (5.2) will follow from

$$1 - \frac{p-1}{2(\alpha+1)p} - 2^{-(p-1)/p} \frac{(\alpha+1)}{2^\alpha} > 0.$$

It is easy to see that when $p \geq 3$, the function $\alpha \mapsto (1+\alpha)2^{-\alpha}$ is an increasing function of α for $0 \leq \alpha \leq 1/p$. It follows from this that for $0 \leq \alpha \leq 1/p$,

$$1 - \frac{p-1}{2(\alpha+1)p} - 2^{-(p-1)/p} \frac{(\alpha+1)}{2^\alpha} > 1 - \frac{p-1}{2p} - 2^{-(p-1)/p} \frac{(1/p+1)}{2^{1/p}} = 0,$$

and from which inequality (5.2) follows.

Now, to show (5.1) holds for all $n \geq 2, p \geq 3$, we first note that for $p > 1$,

$$\sum_{i=1}^n i^{-1/p} < 1 + \int_1^n x^{-1/p} dx = \frac{p}{p-1} n^{1-1/p} - \frac{1}{p-1}.$$

On the other hand, by Hadamard's inequality, which asserts that for a continuous convex function $f(x)$ on $[a, b]$,

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2},$$

we have for $p > 1$,

$$n^{-(p-1)/p} - (n+1)^{-(p-1)/p} = \frac{p-1}{p} \int_n^{n+1} x^{-1-1/p^*} dx \geq \frac{p-1}{p} (n+1/2)^{-1-1/p^*}.$$

Hence inequality (5.1) will follow from the following inequality for $n \geq 2$,

$$\frac{p}{p-1} n^{1-1/p} - \frac{1}{p-1} \leq p^* n^{1/p^*} \left(1 + \frac{1}{2n} \right)^{-(1+p^*)/p}.$$

It is easy to see that for $p > 1$,

$$\left(1 + \frac{1}{2n}\right)^{-(1+p^*)/p} \geq 1 - \frac{1+p^*}{p} \frac{1}{2n}.$$

Hence it suffices to show

$$\frac{p}{p-1} n^{1-1/p} - \frac{1}{p-1} \leq p^* n^{1/p^*} \left(1 - \frac{1+p^*}{p} \frac{1}{2n}\right),$$

or equivalently,

$$\left(1 + \frac{1}{2p-2}\right)^p \leq n.$$

It's easy to check that the right-hand expression above is a decreasing function of $p \geq 3$ and is equal to $5^3/4^3 < 2$ when $p = 3$. Hence it follows that (5.1) holds for all $n \geq 2, p \geq 3$.

We consider lastly inequality (2.4) for other values of α and we take $w_n = n^{\alpha-1/p}$ for $n \geq 1$ so that we can rewrite (2.4) as

$$(5.3) \quad \left(\sum_{i=1}^n i^{\alpha-1/p}\right)^{p-1} < \left(\frac{(\alpha+1)p}{(\alpha+1)p-1}\right)^p \left(\sum_{i=1}^n i^\alpha\right)^p \left(n^{-(p-1)/p-\alpha} - (n+1)^{-(p-1)/p-\alpha}\right).$$

We end our discussion here by considering the case $1 \leq \alpha \leq 1 + 1/p$ and we apply Lemma 2.1 to obtain

$$\begin{aligned} \sum_{i=1}^n i^{\alpha-1/p} &\leq \frac{\alpha-1/p}{\alpha-1/p+1} \frac{n^{\alpha-1/p}(n+1)^{\alpha-1/p}}{(n+1)^{\alpha-1/p} - n^{\alpha-1/p}} = \frac{1}{\alpha-1/p+1} \left(\int_n^{n+1} x^{-\alpha+1/p-1} dx\right)^{-1}, \\ \sum_{i=1}^n i^\alpha &\geq \frac{\alpha}{\alpha+1} \frac{n^\alpha(n+1)^\alpha}{(n+1)^\alpha - n^\alpha} = \frac{1}{\alpha+1} \left(\int_n^{n+1} x^{-\alpha-1} dx\right)^{-1} \end{aligned}$$

We further write

$$n^{-(p-1)/p-\alpha} - (n+1)^{-(p-1)/p-\alpha} = (\alpha-1/p+1) \int_n^{n+1} x^{-\alpha+1/p-2} dx,$$

so that inequality (5.3) will follow from

$$\int_n^{n+1} x^{-\alpha-1} dx < \left(\int_n^{n+1} x^{-\alpha+1/p-1} dx\right)^{1-1/p} \left(\int_n^{n+1} x^{-\alpha+1/p-2} dx\right)^{1/p}.$$

One can easily see that the above inequality holds by Hölder's inequality and it follows that inequality (5.3) holds for $p > 1, 1 \leq \alpha \leq 1 + 1/p$. This provides another proof of inequality (1.3) for $p > 1, 1 \leq \alpha \leq 1 + 1/p$.

REFERENCES

- [1] G. Bennett, Factorizing the classical inequalities, *Mem. Amer. Math. Soc.*, **120** (1996), 1–130.
- [2] G. Bennett, Inequalities complimentary to Hardy, *Quart. J. Math. Oxford Ser. (2)*, **49** (1998), 395–432.
- [3] G. Bennett, Sums of powers and the meaning of l^p , *Houston J. Math.*, **32** (2006), 801–831.
- [4] P. Gao, A note on Hardy-type inequalities, *Proc. Amer. Math. Soc.*, **133** (2005), 1977–1984.
- [5] G. H. Hardy, J. E. Littlewood and G. Pólya, *Inequalities*, Cambridge Univ. Press, 1952.
- [6] K. Knopp, Über Reihen mit positiven Gliedern, *J. London Math. Soc.*, **3** 1928, 205–211 and **5** 1930, 13–21.
- [7] V. I. Levin and S.B. Stečkin, Inequalities, *Amer. Math. Soc. Transl. (2)*, **14** (1960), 1–29.
- [8] R. M. Redheffer, Recurrent inequalities, *Proc. London Math. Soc. (3)*, **17** (1967), 683–699.

DEPARTMENT OF COMPUTER AND MATHEMATICAL SCIENCES, UNIVERSITY OF TORONTO AT SCARBOROUGH, 1265 MILITARY TRAIL, TORONTO ONTARIO, CANADA M1C 1A4

E-mail address: penggao@utsc.utoronto.ca